

	CONTACT	CONTACT PLUS	DROPLET	AIRBORNE
Sign color	Red Sign	Yellow Sign	Green Sign	Blue Sign
Common Diseases	MDRO's: <ul style="list-style-type: none"> • MRSA Active Infection Only • VRE and Hx of VRE • CRE and Hx of CRE • ESBL and Hx of ESBL • Other MDROs • Lice/ Scabies • RSV (<i>if Immunocompromised or Pediatrics</i>) • Shingles localized if unable to cover 	<ul style="list-style-type: none"> • <i>Clostridium difficile</i> • Norovirus 	<ul style="list-style-type: none"> • Bacterial Meningitis or Meningococcemia • Influenza • Pertussis • Rubella • Mumps 	<ul style="list-style-type: none"> • Tuberculosis (TB) or rule-out TB • Varicella (chickenpox) • Shingles Disseminated only (Herpes zoster) • Measles • Novel Airborne Disease (SARS, MERS, etc.)
Alcohol hand gel or hand washing	BEFORE and AFTER patient contact	GEL IN – but AFTER patient/environment contact HAND WASH with soap and water	BEFORE and AFTER patient contact	BEFORE and AFTER patient contact
Private Room *	Yes	Yes	Yes	Yes
Negative Pressure	No	No	No	YES
Closed Door	No	No	No	YES – at all times except to enter/exit room
Mask	No - except as indicated per Standard Precautions	No - except as indicated per Standard Precautions	Yes – surgical or procedure mask	YES (PAPR or N-95)
Gown	Yes	Yes	No - except as indicated per Standard Precautions	No- except as ind per Standard Precautions
Eyewear	Only as indicated per Standard Precautions	Only as indicated per Standard Precautions	Only as indicated per Standard Precautions	Only as indicated per Standard Precautions
Who initiates Isolation Precautions?	<ul style="list-style-type: none"> • MD • Nursing (“within role & scope”) • Infection Control 	<ul style="list-style-type: none"> • MD • Nursing (“within role & scope”) • Infection Control 	<ul style="list-style-type: none"> • MD • Nursing (“within role & scope”) • Infection Control 	<ul style="list-style-type: none"> • MD • Nursing (“within role & scope”) • Infection Control
Who discontinues Isolation Precautions?	<ul style="list-style-type: none"> • ONLY Infection Control or MD 	<ul style="list-style-type: none"> • ONLY Infection Control or MD 	<ul style="list-style-type: none"> • ONLY Infection Control or MD 	<ul style="list-style-type: none"> • ONLY Infection Control or MD • TB – usually requires 3 negative AFB smears or another diagnosis that explains the clinical symptoms
Transport Outside of Isolation Room	<ul style="list-style-type: none"> • Clean transport equipment with BLEACH disinfectant wipes. • Have patient perform hand hygiene • Pt to have clean gown and clean sheet to cover • Transporter /Staff do <u>not</u> wear PPE outside of patient room 	<ul style="list-style-type: none"> • Clean transport equipment with BLEACH wipes. • Have patient perform wash hands with soap and water. • Clean cover gown and clean sheet to cover patient • Transporter /Staff do <u>not</u> wear PPE outside of patient room 	<ul style="list-style-type: none"> • Clean transport equipment with BLEACH disinfectant wipes. • Mask (surgical or procedure) on patient • Transporter/Staff do <u>not</u> wear mask. 	<ul style="list-style-type: none"> • Clean transport equipment with BLEACH disinfectant wipes. • Mask (surgical or procedure) on patient • Transporter/Staff do <u>not</u> wear any type mask or PAPR.
Equipment	<ul style="list-style-type: none"> • Dedicate equipment as possible • Disinfect patient care equipment /high-touch surfaces with hospital-approved BLEACH disinfectant wipes after use. 	<ul style="list-style-type: none"> • Dedicated equipment to room: BP cuff, Thermometer, Stethoscope, WOW • Disinfect patient care equipment / high-touch surfaces with BLEACH wipes after use. 	<ul style="list-style-type: none"> • Dedicate equipment as possible • Disinfect patient care equipment / high-touch surfaces with hospital-approved BLEACH disinfectant wipes after use. 	<ul style="list-style-type: none"> • Dedicate equipment as possible • Disinfect patient care equipment/ high-touch surfaces with hospital-approved BLEACH disinfectant wipes after use.